



LIFE Project Number
LIFE12 ENV/FIN/000409
1st report on the monitoring

Reporting Date
31/03/2015

LIFE+ PROJECT NAME or Acronym

**Climate change indicators and vulnerability of boreal zone applying
innovative observation and modelling techniques**

Project Data

Project location	Helsinki
Project start date:	02/09/2013
Project end date:	01/09/2017 Extension date: <dd/mm/yyyy >
Total Project duration (in months)	48 months (including Extension of <XX> months)
Total budget	2755288 €
Total eligible budget	2755288€
EU contribution:	1366952 €
(%) of total costs	49.61
(%) of eligible costs	49.61

Beneficiary Data

Name Beneficiary	Ilmatieteen laitos
Contact person	Tuula Aalto
Postal address	Erik Palménin aukio 1, FI-00101, Helsinki, Finland
Visit address	Erik Palménin aukio 1, 00560 Helsinki, Finland
Telephone	+358-295395406
Fax:	+358-9-323 2970
E-mail	tuula.aalto@fmi.fi
Project Website	monimet.fmi.fi

MONIMET Action C1

The impact of MONIMET project is monitored with a specific action (C1). The idea was to monitor the impact in the beginning of the project, during the project and at the end of project to see whether there is true communication between the different modeling and experimental approaches and thus convergence of the results at project end. There we defined indicators of the project impact that are common to the different approaches, and thus easily comparable. One such indicator is vegetation active period (VAP), which can be derived from webcam, flux tower, model and remote sensing results. Another indicator is albedo, which can be derived from flux tower, model and remote sensing results. Both VAP and albedo potentially obtain different values depending on the method of estimation, but for example experimental results can be used to tune model parameters, drawing the estimates closer together. In this report we present the methodologies and collect some existing results for VAP and albedo estimation, which can be used in further investigations.

Start of growing season

VAP is the period between the end (GSED) and start day (GSSD) of the growing season. The GSSD is more easily deducible, because in boreal region the springtime development is rapid. Plenty of sunlight is available from early spring, and the clear change from freezing to above zero temperatures driven by synoptic weather patterns initiate the forest carbon exchange. Autumn is more challenging; While the irradiation conditions already start to inhibit photosynthetic uptake, the temperature usually stays at a level where forest could still be active. Furthermore, the dormancy development may initiate before the environmental conditions start to limit carbon exchange. Thus the development towards end of growing season is often gradual and it is difficult to determine a certain ending day. Furthermore, low sun elevation and frequent cloud cover make detection of the GSED difficult from satellite retrievals.

GSSD has been studied by Böttcher et al (2014) using Fractional Snow Cover (FSC), NDVI and NDWI time series from MODIS satellite, phenological observations (Äkäslompolo, Parkano, Paljakka and Saariselkä) and micrometeorological flux measurements (Hyytiälä, Kenttäröva and Sodankylä). Micrometeorological flux measurements and phenological observations were considered as ground truth against which satellite measurements were compared. The GSSD from flux measurements was defined as the day on which the CO₂ uptake exceeds permanently the 15% level of the growing season maximum. The GSSD was obtained from the annual cycle of gross photosynthesis index, which indicates the apparent photosynthetic activity on a daily scale and is related to gross primary production (GPP). It is calculated as a difference of the daily averages of daytime (Photosynthetic Photon Flux Density (PPFD) > 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$) Net Ecosystem Exchange of CO₂ (NEE) and night-time (PPFD < 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$) respiration and is scaled with the growing season maximum. The growing season maximum is the 90th percentile of the daily gross photosynthesis indexes during the month of highest uptake for the multi-year measurement period.

MODIS-derived FSC and vegetation indices (NDVI) were investigated for the use as proxy indicators for GSSD. Daily Terra/MODIS Level-1B data (1 km, 500 m and 250 m products) were collected for the period from February to October 2001–2010. FSC, varying from 0 to 1, was derived at a pixel size of $0.005 \times 0.005^\circ$ using the SCAMod algorithm. The method uses at-satellite

observed single-band reflectance and pixel-wise average forest transmissivity for the provision of FSC.

Vegetation index (NDVI and NDWI) and FSC observations were averaged for all pixels of the seven above mentioned phenology and flux measurement sites. Gaps in daily observations due to cloud cover were filled using linear interpolation and smoothing schemes. GSSD was extracted from NDVI at the time when NDVI's spring-rise began. The general shape of snow depletion curves (changes of FSC during the melt season) for an area can be described by an exponential function, where two parameters are fitted for the melting period. GSSD was derived from the fitted function applying a constant threshold value for FSC.

The satellite indicators detected on average the growing season start 1–2 days later than in situ observations. Correlations between GSSD_NDVI and in situ dates were significant only for site Hyytiälä, whereas significant site-wise correlations between GSSD_FSC and in situ dates were obtained for sites Sodankylä and Hyytiälä, as well as a higher R² for site Kenttäröva. The range of GSSD from the southern to northern sites was captured with the indicators GSSD_NDVI and GSSD_FSC, but data points were grouped for the northern and southern sites, especially for GSSD_NDVI, implying that inter-annual differences at a specific site are less well captured by GSSD_NDVI.

Phenological observations on pine trees at four sites: Parkano, Paljakka, Äkäslompolo and Saariselkä for the years 2001–2010 were obtained from the Finnish Forest Research Institute (METLA). Phenological observations were carried out twice a week in advanced thinning stands. Beginning of shoot elongation can be referred to as growth of pine start date (GPSD). GPSD is usually later than GSSD. MODIS-derived vegetation indices (NDWI) were investigated for the use as proxy indicators for GPSD. GPSD is related to the end of snow melt as described by the decrease of NDWI. Satellite derived GPSD_NDWI accounted only for 54% of the variation in situ observations and data points were dispersed and grouped by site. Site-wise correlations were not significant, hence inter-annual variability at site-level cannot be captured. Mean GPSD for each phenological site could nevertheless be depicted by NDWI timeseries with bias ranging from –2 to 3 days for different sites.

Modeling (JSBACH, the land surface component of MPI-ESM) results have also been compared to remote sensing results. GSSD was extracted from the model annual cycle of gross photosynthesis index similarly to flux observations. The site level correlations were significant for Hyytiälä, but not for Sodankylä. The regional level comparison with satellite results showed that with the multi-year averages the general dynamics of GSSD was rather well captured by JSBACH. However, consistent biases of two to four days were found for GSSD of evergreen conifers when the results were averaged over northern, middle and southern Finland, indicating that the model predicted earlier growing season start than the satellite observation. Biases of three to seven days were found for the deciduous broadleaved forests, indicating later growing season start by the model. There is need for further work to yield better agreement between the different approaches.

The seasonal dynamics of photosynthesis can be related to seasonal variations in ambient temperature. Kolari et al. (2007) derived a sigmoidal relationship between photosynthetic efficiency

and temperature history using in situ observations of pine shoot gas exchange in Värriö and Hyytiälä sites. This result could possibly be used in JSBACH model to tune the photosynthetic potential carboxylation rate (V_{cmax}), in order to make the model respond better to variations in springtime temperatures. In situ fluorescence measurements could also be used in tuning of V_{cmax} . Webcam measurements of the greenness index could also be used in comparisons and model calibration (Fig 1.); when data have been retrieved for more than one year from several sites, there will be enough material for the work.




Fig. 1. Webcam results from Tammela year 2014, showing a distinct springtime increase in the greenness index (gcc).

End of growing season

Though more challenging than start of growing season, procedures have already been developed by the project team for extracting the end of growing season from MODIS satellite NDVI time series and micrometeorological flux measurements (Böttcher et al., 2011). Site comparisons have been made for Hyytiälä, Sodankylä and Kenttäröva. Also phenophase end of height growth of Scots pine (*Pinus sylvestris*) in Parkano and of Norway spruce (*Picea abies*) in Pallasjärvi for years 2001-2010 were utilised.

From flux measurements the following criteria were used for determining the end of season: when daytime Net Ecosystem Exchange (NEE) in autumn first decreases below 20% of the maximum summer uptake (most negative NEE); and when Gross Primary Production (GPP) falls permanently below a certain threshold level of the growing season maximum: 50%, 40%, 30% and 15%. The growing season maximum was defined as the 98th percentile of GPP values during the month of highest uptake for a multi-year period.

The day with maximum NDVI in each year was extracted from linearly interpolated daily MODIS NDVI time-series as indicator for the seasonal vegetation growth peak in coniferous forest. We used the descending inflection point of a double logistic function fitted to the NDVI time-series as satellite-derived end of season indicator for coniferous forest. Furthermore, we evaluated the possibility to use NDVI threshold values for the determination of end of season, scaling NDVI with maximum NDVI.

The 50% GPP indicator showed correspondence with the mid-point of the descending part of the double logistic function fitted to the NDVI time-series at sites Hyytiälä and Kenttäröva for year

2009. The other GPP based phenological indicators occurred later. In Sodankylä flux-based phenological indicators 50% GPP, 40% GPP and 20% NEE, were observed at the same time in year 2010 and seemed to occur when the mid-point of the fitted NDVI profile was already exceeded. Large variations of scaled NDVI for indicators 50% GPP and 40% GPP hindered the application of the threshold for the determination of these events from NDVI time-series.

The day when NDVI had its maximum in summer was compared to phenological observation of end of shoot elongation. The correlation between the two indicators was low. For individual sites, NDVI maximum was observed in average six and two days after end of height growth for Pallasjärvi and Parkano, respectively.

The measure for the end of growing season from NDVI time series revealed low correlations with phenological indicators describing different levels of GPP decrease in autumn at CO₂ flux measurements sites. The data quality of NDVI time-series was in most cases low, especially at the end of autumn, due to low sun elevation and long periods with cloud cover. Furthermore, simple NDVI threshold values seem not to be applicable for extraction of the end of growing season for evergreen coniferous sites. The use of NDVI and other satellite-derived indicators in detecting the end of growing season deserves further studies.

The end of season can be retrieved from the model results using GPP similarly to the procedure described above for the flux measurements. The thresholds applied for the GPP from flux measurements and model results should be re-analysed.

Vegetation active period

The beginning of the season, end of season, and their difference (VAP) will be calculated using the different tools and methods described above. Webcam results will be added to the portfolio. Experimental results will be used to calibrate the models. A first attempt to retrieve VAP from model results is shown below, using a threshold of 15% GPP for both start of season and end of season. The result is averaged over the period from year 1980 to year 2011, and it clearly shows the south-north gradient in VAP.




Fig. 2. Vegetation active period according to JSBACH model results.

Albedo

The total surface albedo can be retrieved from remote sensing as well as models and in situ measurements. The alteration of snow covered and snow free periods create a distinct annual cycle in albedo in boreal region. During the growing season some albedo variation is created through changes in vegetation cover and leaf status. Rinne et al (2007) used NOAA/AVHRR satellite measurements of surface albedo for detecting snow disappearance, which can in turn be connected to start of growing season, as discussed above. Autumn is again more difficult for albedo retrievals.

According to Rinne et al. (2007), the correlation between the snow melt dates based on the remotely-sensed albedo and on local snow depth measurements was reasonably good, and the bias was less than half a day. In comparison to snow depth measurements, the spatial distribution of the timing of the snow melt from albedo observations showed a reasonable pattern with early snow melt in the south and low-lying areas, and late snow melt in the north and high mountains. However, as the data covered only two years, a longer time series is needed to better assess the validity of the method. Here a longer time series could be applied, and compared to snow depth measurements as well as model albedo development and MODIS satellite results.

References

Böttcher K., Aurela M, Markkanen T., Pulliainen J., Metsämäki S., Aalto T., Arslan A.N., Kolari P., Kervinen M., Mattila O.-P. MODIS time-series-derived indicators for the beginning of the growing season in boreal coniferous forest - a comparison with CO₂ flux measurements and phenological observations in Finland. *Remote Sensing of Environment* 140, 625-638, 2014

Böttcher K. et al. 2011 EU-LIFE+ SNOWCARBO (ENV/FIN/000133) Final progress report on extracted features: Action 7: Methodological development and implementation

Kolari P., Lappalainen HK., Hänninen H. and Hari P. Relationship between temperature and the seasonal course of photosynthesis in Scots pine at northern timberline and in southern boreal zone. *Tellus* 50B: 542-552, 2007.

J. Rinne, M. Aurela and T. Manninen. A Simple Method to Determine the Timing of Snow Melt by Remote Sensing with Application to the CO₂ Balances of Northern Mire and Heath Ecosystems. *Remote Sens.* **2009**, *1*, 1097-1107; doi:10.3390/rs1041097